

## Durham Research Online

---

### Deposited in DRO:

05 June 2019

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

McHugh, Cliodhna and Wilson, Fiona and Hind, Karen and Davey, Daniel (2019) 'Cardiovascular health of retired field-based athletes : a systematic review and meta-analysis.', *Orthopaedic journal of sports medicine.*, 7 (8). p. 2325967119862750.

### Further information on publisher's website:

<https://doi.org/10.1177/2325967119862750>

### Publisher's copyright statement:

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License (<http://www.creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

1  
2  
3  
4  
5  
6  
7  
  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

The cardiovascular health of retired field-based athletes: a  
systematic review and meta-analysis.

29   **Abstract**

30   **Background:** Retirement from elite sport participation is associated with decreased physical activity,  
31   depression, obesity and ischemic heart disease. Although engagement in physical activity through  
32   sport is recognised as cardio-protective, an estimated one quarter of American football players  
33   deaths are associated with cardiovascular disease (CVD), predominately players classified as obese.

34   **Hypothesis/Purpose:** To systematically investigate the cardiovascular health profile of retired field-  
35   based athletes.

36   **Methods:** This review was conducted and reported in accordance with the preferred Reporting  
37   Items for Systematic Reviews and Meta-analysis (PRISMA) and pre-registered with PROSPERO. Four  
38   databases (Pubmed, CINAHL, EMBASE, WOS) were systematically searched from inception to  
39   October 2018 using MeSH terms and keywords. Inclusion criteria included; retired field-based  
40   athletes, over the age of 18 years, and at least one CVD risk factor according to the European Society  
41   of Cardiology and American Heart Association. Review articles were not included. Control groups  
42   were not required for inclusion but when available analysis was included. Eligible articles were  
43   extracted using Covidence. Methodological quality was assessed independently by two reviewers  
44   using the AXIS tool. The accuracy of individual study estimates was analysed by random-effect meta-  
45   analyses.

46   **Results:** This review yielded thirteen studies. A total of 4,350 male retired field-based athletes from  
47   two sports; American football and soccer (mean ages: 42.2–66 years) were included. Seven studies  
48   compared retired athletes to control groups. Retired athletes had elevated systolic blood pressure  
49   (BP) in 4 of 6 studies. 50% of studies found greater high-density lipoprotein (HDL), 80% found lower  
50   triglyceride levels and all studies found greater low-density lipoprotein (LDL) for retired athletes  
51   compared to controls. The prevalence and severity of carotid artery calcium (CAC) and carotid artery

52 plaque (CAP) was similar to controls . Retired linemen had double the prevalence of cardio-  
53 metabolic syndrome compared to non-linemen.

54 **Conclusions:** Overall findings are mixed. Inconsistencies in reporting of CVD risk factors and  
55 methodological biases reduced study quality. Retired athletes have comparable CVD risk profile to  
56 the general population. Retired athletes with elevated playing time body mass had an increased  
57 prevalence and severity of risk factors. Significant gaps remain in understanding long term  
58 cardiovascular effects of elite athleticism.

59 **Key Words:** cardiovascular; heart disease; retired athletes; field-based; evidence- based review; risk  
60 factors.

61 **What is known about the subject?** There is no review paper assessing the cardiovascular health  
62 profile of retired field-based athletes. One quarter of retired football player's deaths are caused by  
63 CVD, predominately those classified as obese. There has been a 10% increase in the body mass of  
64 football players over the past decade. Retired athletes often fall into obese classification following  
65 retirement, although it is not known if they experience the same risk factors for CVD as the general  
66 population or if their prior elite athleticism provides a prolonged cardio-protective benefit even in  
67 the presence of obesity.

68 **What this study adds to existing knowledge?** Retired athletes are found to have a substantial risk  
69 for future to adverse CV event, similar to general population indicating athletic past does not  
70 provide long-term protection from CVD. Risk factors associated with CVD; increased systolic BP, LDL,  
71 CAC and cardio-metabolic syndrome are present, particularly those with elevated BMI during playing  
72 career. Conflicting findings are found across studies, highlighting need for further research to clarify  
73 findings.

74

75

## 76 Introduction

77 Regular physical activity is recommended for the optimisation of cardiovascular health and reduction  
78 of all-cause mortality, whereas obesity is an established risk factor for cardiovascular diseases (CVD).  
79 <sup>14, 29, 73, 80</sup> Field-based athletes from sports such as; American football and rugby, present with greater  
80 body mass alongside superior aerobic fitness. <sup>13, 29, 49, 68, 77, 85</sup> However, the cardiovascular health of  
81 these athletes post retirement is unclear. Premature mortality from CVD has become a prominent  
82 topic of discussion for field-based athletes, particularly during the transition into retirement. <sup>4</sup>  
83 Athletes are typically perceived as a healthy cohort, with evidence supporting that fitness provides  
84 protection against known health risks of obesity <sup>14, 41</sup> and relevant co-morbidities. <sup>62, 67</sup> Research also  
85 demonstrates that although exercise has beneficial cardio-protective qualities, it does not  
86 necessarily translate into immunity of cardiovascular risk. <sup>32</sup> An estimated one quarter of American  
87 football players deaths are associated with CVD; predominately those classified as obese.<sup>28</sup>  
88 Furthermore, when compared with retired baseball players, retired football players are more than  
89 twice as likely to die before age 50.<sup>28</sup> The National Institute for Occupational Safety and Health  
90 (NIOSH) study found that although overall mortality from CVD in retired American football players  
91 was 46% lower than the general population, linemen had a 52% greater risk. <sup>4</sup> It has been suggested  
92 that this may in part, reflect increased cardiovascular risk factors associated with greater body size. <sup>4</sup>  
93 <sup>30</sup> The body mass index (BMI) of American football players often falls into the obesity range. <sup>13, 29, 48,</sup>  
94 <sup>77, 85</sup> However, the applicability of BMI in this cohort is widely condemned, due to high muscle mass,  
95 leading to an overestimation of body fat percentage. <sup>55, 56, 58, 70</sup> However, it is unknown if BMI  
96 remains inapplicable to former athletes.

97 To our knowledge, there has been no published review on the evidence concerning  
98 cardiovascular health in retired field-based athletes. The long-term cardiovascular health risks for  
99 professional athletes remain largely unclear. Therefore, the primary purpose of this review was to  
100 systematically collate and appraise the evidence on the cardiovascular health and the prevalence of

101 CVD risk factors in retired field-based athletes. Furthermore, this review aimed to investigate the  
102 prevalence of factors which influence mortality from CVD including: obesity;<sup>33, 36</sup> hypertension;<sup>46</sup>  
103 dyslipidemia;<sup>66</sup> glucose intolerance<sup>20</sup>; cardio-metabolic syndrome (CMS);<sup>20</sup> carotid artery calcium  
104 (CAC)<sup>69, 76</sup> and sleep-disordered breathing (SDB).<sup>64, 74</sup>

## 105 **Methods**

106 This review was conducted and reported in accordance with the Preferred Reporting Items for  
107 Systematic Reviews and Meta-analysis (PRISMA) statement ([www.prisma-statement.org](http://www.prisma-statement.org))<sup>51</sup> and  
108 registered with PROSPERO, a registry of systematic reviews. Registration is available  
109 at <https://www.crd.york.ac.uk/prospero/>; registration number: CRD42017077885. Articles were  
110 retrieved via online database search engines, including; CINAHL, EMBASE, Pubmed, and WOS. The  
111 reference list of all reviews related to cardiovascular health and articles meeting the eligibility  
112 criteria were reviewed manually for suitability.

113 Key words and MeSH terms were searched alone and in combination. The key words  
114 included: CVD, cardiovascular health, blood pressure, lipids, cholesterol, cardio-metabolic syndrome,  
115 hypertension, glucose intolerance, body composition, body mass index, body fat percentage, low-  
116 density lipoprotein, high-density lipoprotein, triglycerides, total cholesterol, sleep-disordered  
117 breathing, field-based athlete, American football, baseball, field hockey, rugby, Gaelic football and  
118 soccer. The inclusion criteria consisted of; retired athletes from a field-based sport, over the age of  
119 18 years and studies that investigated at least one known risk factor for CVD according to the  
120 European Society of Cardiology<sup>62</sup> and the American Heart Association<sup>81</sup>. The search was not limited  
121 by language or publication status and no date restriction implemented. Articles were excluded if the  
122 study design was a review paper and retired athletes with a prior CV event were included. Articles  
123 did not require the inclusion of controls.

**Commented [AK1]:** Query 1: Confirm updates to this sentence per editor's comment.

**Commented [U2R1]:** Confirmed. Amendments were made to this sentence per editor's comments

124 A three-step screening strategy was implemented to identify appropriate relevant articles  
125 for screening using Covidence ([www.covidence.org](http://www.covidence.org)) (Figure 1). Title and abstracts were screened by  
126 two authors blind to each other's selection, in accordance with the aforementioned inclusion  
127 criteria. Then, reviewers independently screened full texts (Appendix 1). A third reviewer was  
128 consulted to make a final decision when consensus was not reached between the two reviewers.  
129 Manual searches were performed of reference lists of the selected articles. The search methodology  
130 and process are described in Figure 1. Eligible articles were critically appraised, and quality assessed  
131 in a full text screening using the AXIS Tool for Critical Appraisal of Cross-sectional Studies (Table 1).<sup>17</sup>  
132 When studies were agreed upon, one reviewer extracted data from selected studies to create an  
133 evidence table, using Strobe guideline analysis.<sup>79</sup>

134 We assessed the precision and accuracy of individual study estimates by implementing  
135 random-effect meta- analyses to examine the overall effect. Heterogeneity between studies was  
136 determined by the  $I^2$  statistic<sup>31</sup> as an indicator of the proportion of total variation in estimates that is  
137 caused by heterogeneity.  $I^2$  values of 25%, 50% and 75% correspond to low, moderate and high  
138 degrees of heterogeneity. Where high levels of heterogeneity were detected ( $I^2 > 75\%$ ) a sensitivity  
139 analysis was implemented. Findings from the random-effects meta-analysis is represented through  
140 forest plots. Studies removed during sensitivity analysis are represented by 0.0% weight in forest  
141 plot figures.




Figure 1: PRISMA Flow Diagram



	Miller <sup>29</sup> (2008)	Panayiotoglou <sup>60</sup> (2017)	Barsa <sup>5</sup> (2014)	Chang <sup>10</sup> (2009)	Hurst <sup>34</sup> (2010)	Hyman <sup>35</sup> (2012)	Albuquerque <sup>1</sup> (2010)	Carruthers <sup>8</sup> (2017)	Kelly <sup>38</sup> (2014)	Lynch <sup>45</sup> (2007)	Pokhare <sup>63</sup> (2014)	Virani <sup>78</sup> (2012)	Luyster <sup>44</sup> (2017)
<b>Introduction</b>													
Were the aims/objectives of the study clear?	N	Y	Y	Y	U	Y	N	N	Y	N	Y	Y	Y
<b>Methods</b>													
Was the study design appropriate for the stated aim(s)?	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Was the sample size justified?	N	N	Y	Y	N	N	Y	N	N	N	N	N	Y
Was the target reference population clearly defined?	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	Y
Was the sample frame taken from an appropriate population base so it closely represented the target/reference population under investigation?	Y	Y	Y	Y	Y	N	Y	U	Y	Y	Y	Y	Y
Was the selection process likely to select subjects/participants that were representative of the target target/reference population under investigation?	Y	Y	Y	Y	Y	Y	U	U	Y	Y	Y	Y	Y
Were there measures undertaken to address and categorise non-responders?	N	N	N	N	N	N	U	N	Y	N	U	U	N
Were the risk factor and outcome variables measured appropriate to the aims of the study?	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Were the risk factor and outcome variables measured correctly using instruments/measurements that had been trialled, piloted or published previously?	Y	Y	Y	Y	Y	U	Y	U	Y	Y	Y	Y	Y
Is it clear what was used to determine statistical significance and/or precision estimates? (e.g. values, CIs)	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	Y	Y

**Commented [AK3]:** Query 2: The journal only prints tables in vertical format. Please check and confirm the changes made to your tables so they are narrower and will fit better on a typeset page.

For Table 1, please confirm edits to the column headings and confirm the abbreviations as added for No, Yes, and Unsure.

**Commented [U4R3]:** Changes to table 1 forma are appropriate and the and abbreviations were confirmed to be correct.

Were the methods (including statistical methods) sufficiently described to enable them to be repeated?	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	Y	Y
<b>Results</b>													
Were the basic data adequately described?	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
Does the response rate raise concern about non-response bias?	Y	Y	N	N	Y	Y	Y	U	N	Y	Y	N	N
If appropriate, was information about non-responders described?	N	N	N	U	U	N	N	U	U	N	U	U	N
Were the results internally consistent?	Y	Y	Y	Y	Y	Y	U	Y	Y	Y	Y	Y	Y
Were the results for the analyses described in methods, presented?	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
<b>Discussion</b>													
Were the authors' discussions and conclusions justified by the results?	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Were the limitations of the study discussed?	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
<b>Other</b>													
Were there any funding sources or conflicts of interest that may affect the authors' interpretation of the results?	N	Y	N	N	N	N	N	N	N	N	N	N	N
Was ethical approval or consent of participants attained?	Y	Y	Y	Y	Y	Y	U	Y	Y	Y	Y	Y	Y
Abbreviations: N, no; Y, yes; U, unsure. Coloured text indicates the following: Green – positive impact on quality of study; Red – negative impact on quality of study; Orange – unknown impact on quality of study													

**Commented [AK5]:** Query 3: state what the different colors for Y, N, and U indicate.

**Commented [U6R5]:** Amended.

## 145   **Results**

146   Overall, 1,816 articles (following the removal of 12 duplicates from the original 1,828 articles) were  
147   retrieved. Screening of titles and abstracts excluded 1,583 articles based on our aforementioned  
148   inclusion criteria. This left 233 articles for full text screening. Of the 233 studies, 76 were excluded as  
149   study participants were current athletes; 132 studies were excluded due to the studies' primary  
150   outcomes not being relevant to traditional cardiovascular health assessment; 10 studies were  
151   excluded due to study design and insufficient data presented and 2 were excluded due to no access  
152   to the full text. Overall, 13 studies were identified as relevant to cardiovascular health from two  
153   field-based sports (American football and Soccer).

154         Of the 13 relevant studies, 11 were cross sectional, 1 observational and 1 prospective study  
155   design. All participants were male and retired professional American football (12/13) or professional  
156   soccer (1/13) players. Twelve studies were conducted in the United States and 1 in Greece (Appendix  
157   1). Studies included consisted of large (n = 948 athletes) and small (n = 12 athletes) cohorts. Studies  
158   compared athletes to age and sex-matched controls, mainly derived from subsets of the following  
159   population cohorts; National Health and Nutrition Examination Survey (NHANES): a longitudinal data  
160   collected between 1999-2006; The Coronary Artery Risk Development in Young Adults (CARDIA):  
161   study of cardiovascular risk development in young adults (n = 5,116); Mayo Clinic Cohort: Mayo  
162   clinic database of all patients who underwent cardiovascular risk evaluation between 2006-2008);  
163   Dallas Heart Study (DHS): probability based cohort of Dallas County adults, oversampled for African-  
164   Americans (n = 6,101) and The Aerobics Centre Longitudinal Study (ACLS): longitudinal study of  
165   medical health. Participants were selected from a subset of 5,322 of the total 17,967 participants.  
166   (Appendix 1). Analysis was carried out to assess CVD risk factors according to the ESC and AHA risk  
167   factor guidelines, to compare risk factors based on sport, playing position to control groups. The  
168   studies included evaluated cardiovascular health and CVD risk in retired athletes under the following  
169   categories (Table 3); Body Composition: BMI, body fat percentage, waist circumference, neck

170 circumference, waist to hip ratio; Blood pressure (BP); Lipids: total cholesterol, high-density  
 171 lipoprotein (HDL), low-density lipoprotein (LDL), triglycerides, total cholesterol/HDL,  
 172 triglycerides/HDL; Glucose; Cardio-Metabolic Syndrome; <sup>27</sup> Carotid Artery Calcium (CAC/Carotid  
 173 Artery Plaque (CAP); <sup>26</sup> Sleep Disordered Breathing (SDB). <sup>72</sup>  
 174  
 175 Body Composition:  
 176 Ten studies assessed BMI, accounting for 3,095 retired athletes (8-American football, 1-Soccer) with  
 177 a mean BMI of 29.3 kg/m<sup>2</sup> (American football: 31.3 kg/m<sup>2</sup>; Soccer: 27.3 kg/m<sup>2</sup>). American football  
 178 players had greater mean BMI than controls. <sup>1, 8, 10, 44</sup> Compared to controls, retired athletes had  
 179 lower body fat percentages. <sup>45, 60</sup> Lower waist circumference values were reported for retired athletes  
 180 compared to the Dallas DHS, CARDIA and a control group. <sup>10, 44, 45</sup> However, compared to ACLS retired  
 181 footballers had a higher mean waist circumference (Table 2). <sup>10</sup> Conflicting waist-to-hip ratio  
 182 findings were reported for retired American football athletes <sup>10, 45</sup> Waist-to-hip ratio was found to be  
 183 considerably lower for retired soccer players than values for American football players. Using BMI ≥  
 184 30kg/m<sup>2</sup>, 67.4% of retired players were classified as obese compared to 10% when using BF% ≥ 25%  
 185 via dual-energy X-ray absorptiometry (DXA). <sup>35</sup>  
 186 Sub-group analysis found that retired linemen had elevated measures of body composition  
 187 compared to retired non-linemen; higher BMI <sup>5, 34, 44, 50</sup> and higher body fat percentages. <sup>50</sup>  
 188 Conflicting findings were reported on waist circumference between retired linemen and retired non-  
 189 linemen. <sup>5, 44</sup>

Table 2: Body Composition Measures

Lead Author	BMI (kg/m <sup>2</sup> )	WC/NC (cm)	BF (%)	WHR
Miller <sup>50</sup>	LM v NLM: 34.9 (4.9) v 30.7 (4.0), ***		LM v NLM: 31.4% v 27.4%, *** BF% >28%: LM v NLM: 111 (67.7%) v 145 (41.9%), ***	
Panayiotoglou <sup>60</sup>	Soccer v Controls: 27.3 ± 2.8 v 27.4 ± 2.7, ns p = 0.919		Soccer v Controls: 24.5 ± 4.5 v 27 ± 3.9, ns	Soccer v Controls: 0.96 ± 0.05 v 0.97 ± 0.01, ns
Barsa <sup>5</sup>	LM v NLM: 33.6 (30.5-37.9) v 30.3 (27.7-33), p < 0.001	LM v NLM: 109.2 (99.1-119.4) v 99.1 (91.4-106.6), p < 0.001		
Chang <sup>10</sup>	NFL v DHS: 31.5 ± 4.2 v 31.4 ± 4.0, ns NFL v ACLS: 31.7 ± 4.7 v 28.6 ± 3.1, ***	NFL v DHS: 103.8 ± 11.5 v 107.4 ± 10.9, ns NFL v ACLS: 105.7 ± 12.7 v 98.4 ± 8.9, ***		NFL v DHS: 1.08 ± 0.85 v 0.98 ± 0.05, *** NFL v ACLS: 1.06 ± 0.73 v 0.93 ± 0.05, ***
Hurst <sup>34</sup>	NFL v Mayo: 31.5 ± 4.5 v 31.0 ± 2.7, ns LM v NLM: 34.2 ± 4.5 v 30.5 ± 4.0, ***			
Hyman <sup>35</sup>	BMI >30 = 89 (67%) BMI correlated with LM ***	BF% > 25% =13 (10%)		
Albuquerque <sup>1</sup>	NFL v Controls: 32.3 ± 0.3 v 30.0 ± 0.1. p < 0.001			
Carruthers <sup>8</sup>	NFL v DHS: 32.5 (5.4) v 29.3 (5.4), ***			
Kelly <sup>38</sup>	BMI: 33.8 ± 6 BMI >30: 45 (66.2%)			
Lynch <sup>45</sup>	NFL v Controls: 29.4 ± 2.8 v 30 ± 3, ns	NFL v Controls: 101.2 ± 6.8 v 106.1 ± 8.0, ns	NFL v Controls: 23 ± 4 v 32 ± 7, ***	NFL v Controls: 0.95 ± 0.05 v 0.98 ± 0.06, ns
Pokharel <sup>63</sup>	BMI: 31 (29-35)	WC: 40 (37-44) NC: 17 (16-18)		

**Commented [AK7]:** Query 4: Confirm the edits to the leftmost column for conciseness. Make any further edits to reduce the overall width of the table.

**Commented [U8R7]:** Adjustments made to table to reduce overall width of table

**Commented [AK9]:** Query 5: suggest nonsignificant *P* values be listed simply as “ns” for conciseness, throughout all tables.

**Commented [U10R9]:** Amended

Virani <sup>78</sup>		WC: 39.4 ± 10.6
Luyster <sup>44</sup>	NFL v CARDIA: 30.3 ± 3.8 v 29.9 ± 4, ns	NFL v CARDIA: 95.2 ± 22 v 98.1 ± 10.2, ns
	LM v NLM: 29.8 ± 2.9 v 30.5 ± 4.1, ns	LM v NLM: 92.6 ± 19.7 v 96.2 ± 22.8, ns
	BMI ≥ 30: NFL v CARDIA: 60 (49.2%) v 51 (41.8%), ns	
	LM v NLM: 13 (38.2%) v 47 (53.4%), ns	

191 Abbreviations: LM: linemen; NLM: non-linemen; BMI: body mass index; WC: waist circumference; NC: neck circumference; BF%: body fat percentage; WHR:  
192 waist-to-hip ratio; NHANES: National Health and Nutrition Examination Survey; CARDIA: The Coronary Artery Risk Development in Young Adults Mayo;  
193 Mayo Clinic Cohort; DHS: Dallas Heart Study; ACLS: The Aerobics Centre Longitudinal Study; NFL: National Football League.

194 Data are reported as: mean ± SD, median (IQR) or n (%). ns – non-significant; \* - p<0.05; \*\* - p<0.01; \*\*\* - p<0.001.

**Commented [AK11]:** Query 6: List in a general footnote how the data in the table are reported: mean ± SD, n (%), etc. It's unclear how the data are presented for Carruthers BMI, Barsa WC/NC, and Pokharel WC/NC

**Commented [U12R11]:** Amended

195    *Hypertension and Blood Pressure:*

196    A greater incidence of hypertension was reported in athletes compared to controls.<sup>1, 5, 34, 44</sup> Retired  
197    linemen had higher prevalence of hypertension compared to retired non-linemen, although not  
198    significantly (Table 3).<sup>5, 50 34, 44</sup> Most studies found higher resting systolic BP in retired football  
199    players compared to controls (Table 3).<sup>1, 8, 10, 34, 44, 45</sup> Conflicting findings were reported for sub-group  
200    analysis in American football players. (Table 3)<sup>5, 34, 44, 50</sup>

201

202    *Lipid Profiles:*

203    Eleven studies analysed measures of lipid profiles. A greater prevalence of hyperlipidemia was  
204    reported for retired athletes compared to controls and population norms.<sup>10, 34</sup> Average total  
205    cholesterol for retired athletes was 194.3 mg.dl. Mixed findings were reported for mean *total*  
206    *cholesterol* for retired athletes; three studies report lower values<sup>1, 10, 34</sup> and three report higher  
207    values (Table 3).<sup>10, 44, 45</sup> Most studies examining retired football athletes found higher HDL values  
208    compared to controls.<sup>1, 8, 10, 34, 44, 45</sup> Mean LDL for retired American football and soccer players was  
209    123.9 mg.dl and 134mg.dl, respectively, both above recommended levels.<sup>1, 5, 10, 25, 34, 44, 45, 63, 78</sup> All  
210    studies reported higher LDL values for retired athletes compared to controls.<sup>1, 10, 34, 44, 45</sup> Five of six  
211    studies reported lower triglyceride values in retired athletes.<sup>1, 10, 34, 45, 60</sup> One of three studies  
212    investigating total cholesterol/HDL ratio reported lower ratios in retired athletes compared to  
213    controls.<sup>34, 45, 60</sup>

214            Sub-group analysis based on prior playing position for American football showed that retired  
215    linemen had higher total cholesterol values in three studies<sup>5, 34, 50</sup> and equal in one.<sup>44</sup> Four studies  
216    found higher HDL values in retired compared to non-linemen.<sup>5, 34, 44, 50</sup> Conflicting findings for LDL  
217    were reported; two reported higher values for non-linemen<sup>5, 34</sup> and one showed higher for linemen

218 (Table 3).<sup>44</sup> Inconsistencies were found triglyceride levels; Two studies reported higher levels for  
219 linemen<sup>34, 50</sup> and two reported higher levels for non-linemen.<sup>5, 44</sup>

220 *Glucose:*

221 The prevalence of diabetes was reported to be between 7% and 8%.<sup>1, 5, 10, 35, 50, 63, 78</sup>

222 Conflicting reports on fasting glucose was reported for retired football players; three studies found  
223 lower glucose values for retired football players and two found higher levels compared to  
224 comparators.<sup>1, 10, 34, 44, 45</sup> Higher glucose values were reported for former linemen compared to non-  
225 linemen.<sup>5, 34, 44, 50</sup>

226 *Biomarkers:*

227 Four studies measured high sensitive C-reactive protein (hsCRP.)<sup>5, 10, 63, 78</sup> Chang reported  
228 conflicting findings; lower values in retired athletes compared to DHS but higher values compared to  
229 ACLS.<sup>10</sup> Two studies found no association between hsCRP levels and CAP or subclinical  
230 atherosclerosis.<sup>5, 78</sup> HsCRP was found to be significantly higher in retired NFL players with pre-  
231 existing cardio-metabolic syndrome.<sup>78</sup>

232



**Table 3: Risk Factors for CVD**

Lead Author	Blood Pressure	Lipids	Carotid Artery Calcium	Cardio-Metabolic Syndrome
Miller <sup>50</sup>	<b>LM v NLM</b> HT: 41 (25%) v 71 (20.5%), ns SBP: 137.1 (21.3) v 131.9 (17.4), ** DBP: 79.2 (13.3) v 78.5 (11.4), ns	<b>LM v NLM:</b> HDL: 44.5 (14.2) v 47.6 (14.9), * TC: 189.1 (43.9) v 195.6 (38.6), ns TG: 128.5 (79.8) v 116.1 (70.8), ns		<b>LM v NLM:</b> CMS: 98 (59.8%) v 104 (30.1%), *** BMI $\geq 30$ kg/m <sup>2</sup> : 140 (85.4%) v 174 (50.3%), *** Raised BP: 111 (67.7%) v 212 (61.3%), ns Reduced HDL: 69 (42.1%) v 113 (32.7%), ns Raised FG: 99 (60.4%) v 130 (37.6%), *** Raised TG: 51 (31.1%) v 83 (24%), ns
Panayiotoglou <sup>60</sup>		<b>Soccer v Controls:</b> TG: $1.1 \pm 0.2$ v $1.6 \pm 0.8$ , ns		
Barsa <sup>5</sup>	<b>LM v NLM:</b> HT: 38.8% v 28.5%, ** SBP: 131 (122-144) v 130 (120-143), ns	<b>LM v NLM:</b> HDL: 45 (39-55.8) v 48 (40-57), ** LDL: 117.5 (98-143) v 127 (104-151.8), *** TC: 190 (167.5-214) v 198 (173-227), * TG: 88 (62-141) v 91 (66-140.5), ns	<b>LM v NLM:</b> CAC = 0: 105 (33.88%) v 259 (41.7%), * CAC = 1-100: 103 (33.22%) v 198 (31.88%), ns CAC $\geq 100$ : 102 (32.9%) v 164 (26.41%), *	<b>LM v NLM:</b> CMS: 25.8% v 16.5%, ***
Chang <sup>10</sup>	<b>NFL v DHS:</b> SBP: $127.6 \pm 16.7$ v $135.6 \pm 17.0$ , *** DBP: $77.3 \pm 11.2$ v $82.5 \pm 10.4$ , p *** <b>NFL v ACLS:</b> SBP: $129.2 \pm 17.0$ v $129 \pm 16$ , ns DBP: $77.5 \pm 11.1$ v $85.0 \pm 9.8$ , ***	<b>NFL v DHS:</b> HDL: $50.8 \pm 16.8$ v $43.7 \pm 10.9$ , *** LDL: $128.5 \pm 36.0$ v $107.7 \pm 37.5$ , p *** TC: $197.8 \pm 42.1$ v $176.8 \pm 40.1$ , *** TG: 81 (61-115) v 111 (74-160), *** <b>NFL v ACLS:</b> HDL: $49.4 \pm 17.0$ v $46.4 \pm 11.5$ , ns	<b>NFL v DHS:</b> 46% v 48.3%, ns  No statistically significant difference across CAC scores for all groups	<b>NFL v DHS:</b> Significantly lower percentage of retired players with CMS compared to controls *  <b>NFL v ACLS:</b> 39.5 % v 23%, ***

**Commented [AK13]:** Query 7: Confirm the edits to this column for conciseness. Make any further edits to reduce the overall width of the table.

**Commented [U14R13]:** Edits to the table were checked and clarified.

**Commented [U15R13]:** Amendments made to table to reduce overall width

		LDL: 126 ± 36.2 v 124.7 ± 37.2, ns TC: 192.9 ± 41.9 v 204.0 ± 41.6, *** TG: 83.5 (61-122) v 127.5 (92-177), ***		
Hurst <sup>34</sup>	<b>NFL v Mayo:</b> HT: 38(19%) v 6(7%). SBP: 128.7 ± 16.4 v 123.7 ± 13.8, ** DBP: 78.7 ± 10.9 v 78.4 ± 8.2, ns <b>LM v NLM:</b> HT: 12(20%) v 9(6%), * SBP: 128.8 ± 16.9 v 128.6 ± 16.2, ns DBP: 79.2 ± 13.1 v 78.6 ± 9.9, ns	<b>NFL v Mayo:</b> HDL: 40.9 ± 16.5 v 50.1 ± 13.5, ns LDL: 131.3 ± 25.6 v 126.4 ± 35.5, ns TC: 198.8 ± 40.8 v 207.2 ± 40.1, ns TG: 102.6 ± 64.6 v 162.2 ± 128.3, *** <b>LM v NLM:</b> HDL: 45.4 ± 18.4 v 50.5 ± 15.6, p = 0.045 LDL: 127.5 ± 30.1 v 132.8 ± 37.7, ns TC: 197.7 ± 37 v 199.2 ± 42.5, ns TG: 120.5 ± 64.4 v 95.1 ± 63.4, **	<b>NFL v Mayo:</b> Plaque: 67 (33%) v 36 (29%), ns  <b>LM v NLM:</b> Plaque: 16(27%) v 51(36%), ns	<b>LM v NLM:</b> 27(46%) v 32 (23%), ***
Hyman <sup>35</sup>	HT: 55 (42.6%)			
Albuquerque <sup>1</sup>	<b>NFL Vs Controls:</b> HT: 37.8% v 21.4%, *** SBP: 133.5 ± 1.1 v 126.5 ± 0.5, *** DBP: 80.0 ± 0.7 v 72.7 ± 0.3, ***	<b>NFL Vs Controls:</b> HDL: 44 ± 0.8 v 47 ± 0.3, *** LDL: 121.4 ± 2.3 v 117 ± 1.3, ns TC: 183.4 ± 4.1 v 195.3 ± 1.5, ns TG: 149.8 ± 12.7 v 168 ± 4.7, ***		
Carruthers <sup>8</sup>	<b>NFL v DHS:</b> SBP: 136.1 (17.2) v 132.7 (17), *	<b>NFL v DHS:</b> HDL: 55.9 (16.6) v 48.9 (12.9), ***	<b>NFL v DHS:</b> <b>Median (95% CI) CAC:</b> 0.5 (0, 45.2) v 1.8 (0, 73), ns <b>&lt;5% risk:</b> CAC = 0: 23 (60%) v 120 (64%), ns CAC = 1-100: 12 (32%) v 56 (30%) CAC ≥100: 3 (8%) v 11 (6%) <b>5-7.5% Risk:</b> CAC=0: 15 (65%) v 61 (48%), * CAC=1-100: 3 (13%) v 53 (41%) CAC ≥100: 5 (22%) 14 (11%) <b>&gt;7.5% Risk:</b> CAC = 0: 14 (32%) v 83 (27%), **	

Commented [AK16]: Query 8: median (95% CI)?

Commented [U17R16]: Included

			CAC = 1-100: 21 (49%) v 116 (38%) CAC ≥100: 8 (19%) v 104 (34%)	
Kelly <sup>38</sup>				34 (50%) 234
Lynch <sup>45</sup>	<b>NFL v Controls:</b> SBP: 130 ± 19 v 133 ± 20, ns DBP: 79 ± 8 v 82 ± 13, ns	<b>NFL v Controls:</b> HDL (mM): 1.30 ± 0.23 v 0.95 ± 0.19, *** LDL (mM): 3.10 ± 0.48 v 3.04 ± 0.61, ns TC (mM): 4.93 ± 0.52 v 4.75 ± 0.76, ns TG (mM): 1.17 ± 0.69 v 1.71 ± 0.67, *		
Pokharel <sup>63</sup>	HT: 267 (32%) SBP: 130 (121-142) DBP: 80 (74-87)	HDL: 47 (39-56) LDL: 125 (103-148) TC: 196 (171-223) TG: 89 (64-141)		239
Virani <sup>78</sup>	HT: 309 (34.7%)	HDL: 49 ± 14 LDL: 127 ± 38 TC: 199 ± 41 TG: 89 ± 77	CAC detected in 41% of players	187 (19.7%) 240
Luyster <sup>44</sup>	<b>NFL Vs CARDIA:</b> HT: 36 (29.5%) v 35 (28.7%), ns SBP: 125.3 ± 13.9 v 120.4 ± 13.2, ** DBP: 80.1 ± 10.3 v 75 ± 11.2, *** <b>LM v NLM:</b> HT: 9 (26.5%) v 27 (30.7%), ns SBP: 124.6 ± 16.9 v 125.6 ± 12.7, ns DBP: 81.4 ± 9.7 v 80 ± 9.7, ns	<b>NFL Vs CARDIA:</b> HDL: 49.9 ± 11.5 v 44.9 ± 12, *** LDL: 126.5 ± 39.7 v 110.5 ± 31.7, *** TC: 197.9 ± 43.5 v 183.4 ± 35.9, ** TG: 140.3 ± 96.5 v 140.2 ± 92.4, * <b>LM v NLM:</b> HDL: 47.9 ± 11.9 v 50.8 ± 11.3, ns LDL: 135 ± 48.7 v 122.9 ± 34.9, ns TC: 198.1 ± 55.8 v 197.9 ± 37.6, ns TG: 105.3 ± 75.1 v 120.5 ± 104.3, ns	<b>NFL v CARDIA:</b> CAC Presence: 37 (30.3%) v 37 (30.3%), ns CAC Distribution: ns CAC = 0: 87 (71.3%) v 87 (71.3%) CAC = 1-99.99: 29 (23.8%) v 28(23%) CAC ≥100: 6 (4.9%) v 7 (5.7%). <b>LM v NLM:</b> CAC presence: 8 (23.5%) v 29 (33%), ns	242 243 244 245 246 247

248 Abbreviations: LM: linemen; NLM: non-linemen; NHANES: National Health and Nutrition Examination Survey; CARDIA: The Coronary Artery Risk  
249 Development in Young Adults Mayo; Mayo Clinic Cohort; DHS: Dallas Heart Study; ACLS: The Aerobics Centre Longitudinal Study; NFL: National Football  
250 League; SBP: systolic blood pressure; DBP: diastolic blood pressure; HDL: high-density lipoprotein; LDL: low-density lipoprotein; TC: total cholesterol; TG:

251 triglycerides; CAC: carotid artery calcium; CAP: carotid artery plaque; IFG: impaired fasting glucose; CMS: cardio-metabolic syndrome. Data are reported as:  
252 mean ± SD, median (IQR) or n (%). ns – non-significant; \* - p<0.05; \*\*p<0.01; \*\*\* - p<0.001

**Commented [AK18]:** Query 9: List in a general footnote how the data in the table are reported: mean ± SD, median (range), n (%), etc.

**Commented [U19R18]:** Amended

**Commented [AK20]:** Query 6: List in a general footnote how the data in the table are reported: mean ± SD, n (%), etc. It's unclear how the data are presented for Carruthers BMI, Barsa WC/NC, and Pokharel WC/NC

**Commented [U21R20]:** Amended

253 *Cardio-metabolic Syndrome:*

254 Six studies reported on the prevalence of cardio-metabolic syndrome in retired American football  
255 players. A substantial variance was found, ranging from 19.7% to 50%. Compared to the DHS a  
256 significantly lower incidence of cardio-metabolic syndrome was reported; however, compared to  
257 ACLS and controls a higher prevalence was reported for retired football players.<sup>5, 10</sup> Retired linemen  
258 had almost double the prevalence of cardio-metabolic syndrome compared to non-linemen (Table  
259 3).<sup>5, 34, 50</sup> Three of the component criteria; BMI  $\geq 30\text{kg.m}^2$ , reduced HDL cholesterol and raised  
260 fasting glucose, were significantly greater in retired linemen compared with non-linemen.<sup>50</sup>

261 *Carotid Artery Calcium/Carotid Artery Plaque:*

262 Three studies found similar prevalence and severity of CAC between retired athletes and controls.<sup>8,</sup>  
263 <sup>10, 44</sup> When controlled for ethnicity, no difference in CAC between retired players and DHS  
264 (Caucasians: 67.2% v 57.4%; African Americans: 31.5% v 42.1%) was reported.<sup>10</sup> Conflicting findings  
265 were reported for sub-group analysis of American football players. Two studies reported a greater  
266 prevalence of CAC/CAP for non-linemen.<sup>34, 44</sup> However, Miller reported that retired linemen were  
267 less likely to have absence of CAC, a similar likelihood of mild CAC and a greater likelihood of  
268 moderate to severe CAC compared to non-linemen (Table 3). Furthermore, adjusting for  
269 demographic and metabolic covariates, linemen playing position remained independently associated  
270 with mild (OR 1.41; 95% CI: 1.05, 2.2) and moderate to severe (OR 1.67; 95% CI: 1.05, 2.2) subclinical  
271 atherosclerosis.<sup>5</sup>

272 *Sleep-disordered Breathing:*

273 A limited number of studies analysed SDB. Self-reported presence of obstructive sleep  
274 apnoea in retired football players was reported between 41%-53%.<sup>1 35</sup> One study reported retired  
275 football players had double the prevalence of high risk sleep apnoea compared to the CARDIA study.  
276 <sup>44</sup> Retired soccer players reported less days snoring per week than controls.<sup>60</sup>

277 *Smoking:*

278 Retired athletes were reported to have lower prevalence of smoking (past or present), in all  
279 studies<sup>1, 8, 10, 34, 44, 50, 60</sup> Conflicting findings were identified based on playing position in American  
280 football. Two studies reported lower prevalence of smoking history in linemen compared to non-  
281 linemen,<sup>44, 50</sup> whereas, one study reported a greater prevalence in linemen.<sup>34</sup>

282


283 *Meta-analysis:*

284 Implementation of meta-analysis using random-effects and sensitivity analysis indicated that the  
285 overall effect of prior engagement in American football had a positive effect on fasting glucose  
286 levels, finding a mean difference of -4.66 (95%CI: -7.71, -1.62,  $I^2=55\%$ ) when compared to controls  
287 (Figure 2). Prior engagement in American football had a negative effect on systolic BP, with a mean  
288 difference of 3.07 (95%CI: 0.78, 5.36,  $I^2=44\%$ ) in favour of controls (Figure 3). A wide confidence  
289 interval was identified for triglycerides of athletes. A mean difference of -19.07 (95%CI: -34.96, -3.19,  
290  $I^2=59\%$ ) in favour of retired athletes was found for triglycerides (Figure 4). Retired players had a  
291 higher mean value for LDL compared to control groups with a mean difference of 5.00 (95%CI: 1.54,  
292 8.47,  $I^2=42\%$ ) (Figure 5).

293 *Risk of Bias:*

294 Studies were critically appraised using the AXIS tool.<sup>17</sup> Overall, studies were of moderate quality  
295 with common issues identified in several domains. Where 'unsure' response was assigned it was  
296 most commonly associated with lack of clarity in reporting. Many studies did not provide  
297 justification for the sample size due to their cross-sectional and observational study design. Studies  
298 did not address the issue of non-responders; provide information or categorise. Samples of  
299 convenience were most commonly sought, and it was not addressed how representative these  
300 samples were to the true population.

301



302

Figure 2: Forest Plot of Systolic BP.

303





Figure 3: Forest Plot of Glucose

304



305

Figure 4: Forest Plot of Triglycerides.

306




Figure 5: Forest Plot of LDL.



## 311 Discussion

312 Thirteen studies examined the cardiovascular health profile of retired field-based athletes. The  
313 variance in study objectives provides a broad understanding of the cardiovascular health and CVD  
314 risk profile of retired contact sport athletes and how this compares to the general population.

315         Synthesis of studies suggests that retired athletes with elevated BMI have a similar risk for  
316 future adverse cardiovascular event as obese non-athletes from the general population. The  
317 Framingham study in 2008 indicated that men and women who are obese have a lifetime risk of CVD  
318 of 66.8% and 46.7%, respectively.<sup>21</sup> Obesity measured by BMI is common among retired football  
319 players and found to be more prevalent in 5 of 6 studies (83%) compared to comparators.<sup>1, 8, 10, 34, 44</sup>  
320 There is an argument that because BMI doesn't consider the increased muscle mass in current  
321 athletes, as a measure of body composition, BMI may overestimate the prevalence of obesity in  
322 current athletes. Despite a threefold greater engagement in physical activity during young adulthood  
323 (20-34 years), after the age of 65, former athletes (Age (yr):  $66 \pm 6$ ) and sedentary individuals (Age  
324 (yr):  $67 \pm 5$ ) have similar levels of physical activity.<sup>45</sup> The errors associated with using BMI during  
325 active athletic career are most likely not as significant during retirement, therefore using BMI as a  
326 measure of obesity during retirement is more appropriate Epidemiological research has consistently  
327 reported increased risk of cardiovascular death with increased BMI.<sup>15, 40, 59, 65</sup> Retired linemen were  
328 found to have elevated measures of body composition compared to non-linemen in all but one  
329 study.<sup>5, 34, 44, 50</sup> Luyster suggested that an average of 25 years following retirement, non-linemen  
330 have equal probability of becoming obese as linemen. Similarly, Miller reported that 50% of retired  
331 non-linemen had a BMI  $\geq 30\text{kg/m}^2$ .<sup>44, 50</sup> The average age of retired NFL players was 57.1, falling in  
332 line with the estimated 37.7% prevalence of obesity (BMI  $\geq 30\text{kg/m}^2$ ) in males between the ages of  
333 40-59 from the general population.<sup>22</sup> Despite known limitations, BMI remains the most widely used  
334 measure of obesity, with 76.9% (10/13) of studies in this review applying it. It is postulated that  
335 waist circumference and waist-to-hip ratio are more accurate indicators of obesity and future risk of

336 CVD for athletes than BMI. <sup>9, 47</sup> Interestingly, when matched for BMI both Luyster and Chang (DHS  
337 cohort) reported lower waist circumference levels. <sup>10, 44</sup>

338 This highlights the need for more reliable measures of body composition beyond BMI, waist  
339 circumference and waist-to-hip ratio, such as DXA scans. DXA scans provide in-depth analysis of  
340 body composition, identifying, lean mass, fat mass and visceral adipose tissue (VAT). In current field-  
341 based athletes, elevated BMI often reflects greater lean muscle mass. <sup>3, 24, 37, 42, 52</sup> Findings from this  
342 review suggests this may not be the case in retired players. This generates speculation that  
343 persistent reporting of elevated BMI found in retired athletes reflects an increase in fat mass,  
344 compared to the increased lean mass found in current athletes. No study analysed VAT in this  
345 review. Epidemiological research has consistently reported links between VAT and systemic  
346 inflammation. <sup>19, 61, 71, 86</sup> It remains unclear if elevated BMI during retirement diminishes benefits  
347 gained from an individual's past elite athleticism.

348 The cause of long-term risk of elevated BP and hypertension in retired-field based athletes is  
349 unclear. In current American football athletes, there is an increased prevalence of hypertension  
350 (13.8%) compared to age-and-sex matched controls (5.5%). <sup>77</sup> The 1994 study by NIOSH reported;  
351 deaths among linemen were almost exclusively attributable to hypertension and ischemic heart  
352 disease. <sup>4</sup> According to the American Heart Association, risk of death from ischemic heart disease  
353 and stroke doubles with every 20 mm Hg systolic or 10 mm Hg diastolic increase among people from  
354 age 40 to 89. <sup>81</sup> Average systolic BP was 130.6mm Hg in this review, 10mm Hg above recommended  
355 target levels. <sup>7, 81</sup>

356 Where elevated BP was reported concomitant increases in body composition were typically  
357 reported. <sup>1, 8, 34</sup> When matched for BMI results on BP are conflicting. Compared to DHS cohort,  
358 retired football players had lower BP <sup>10</sup>, however Hurst and Luyster reported higher BP. <sup>34, 44</sup> No  
359 study controlled for smoking, alcohol intake or dietary intake, therefore it's not possible to identify

**Commented [AK22]:** Query 10: Confirm this change, per editor's comment.

**Commented [U23R22]:** Change to adipose confirmed as per editor's comments.

360 the cause of higher mean BP in retired players. Furthermore , studies comparing retired NFL players  
361 based on positions reported similar or increased BP for linemen <sup>5, 34, 44, 50</sup>

362 This suggests it's possible that body composition in linemen might offset some benefits of  
363 exercise on BP. Retired soccer players mean BMI was  $\leq 30\text{kg.m}^2$  ( $27.2\text{kg.m}^2$ ), however according to  
364 the European Society of Hypertension Guidelines,<sup>18</sup> 66% had BP exceeding the upper range of grade  
365 1 hypertension. This corresponds to statistics from age-sex matched individuals from the general  
366 population.<sup>82</sup> While overall non-linemen had statistically similar BMI to linemen ( $29.8\text{ kg.m}^2$  v  $30.5$   
367  $\text{kg.m}^2$ ) but a greater percentage of African-American athletes ( $26.4\%$  v  $47.7\%$ ,  $p = 0.03$ ), the  
368 incidence of hypertension was higher for non-linemen ( $26.5\%$  v  $30.7\%$ ).<sup>44</sup> Meta-analysis identified a  
369 moderate statistical heterogeneity ( $I^2=44\%$ ) (Figure 3). A possible cause for the high level of  
370 heterogeneity in these studies is the significant difference in smoking history, and BMI between  
371 retired players and controls, along with measures that were not controlled for.

372 An increased prevalence of hyperlipidemia and fasting glucose in retired linemen was  
373 reported in 2 studies.<sup>10, 34</sup> Results were conflicting for LDL and HDL concentrations. <sup>1, 8, 10, 34, 44, 45, 60</sup>  
374 Higher levels of HDL in retired athletes coincided with higher levels of total cholesterol <sup>8, 10, 44, 45</sup> This  
375 may be attributed to their physically active and high caloric diet past. Where lower levels of HDL  
376 were reported in retired players elevated measures of body composition was reported  
377 (Albuquerque; Hurst). . This suggests that size matters and early cardiovascular risk factor screening  
378 and maintenance of physical activity levels in early retirement is needed. <sup>1, 10, 34, 50</sup>. Multiple papers  
379 reported that hyperlipidemia is associated with coronary atherosclerosis, CAP and cardio-metabolic  
380 syndrome. <sup>10, 45, 50, 78</sup> All studies reported that retired players had higher LDL concentration compared  
381 to controls; this may be caused by high caloric diet during career or the change in body composition  
382 that occurs during retirement. <sup>1, 10, 34, 44, 45</sup> Increasing BF% coincides with a decrease in physical  
383 activity levels, similar to that of obese sedentary controls after the age of 65 years. <sup>45</sup>

384 A lower prevalence of diabetes was reported for retired athletes, despite indication of a  
 385 threefold higher prevalence of impaired fasting glucose.<sup>1, 10, 44</sup> When stratified by ethnicity, African-  
 386 American athletes had a significantly greater prevalence of impaired fasting glucose, whereas,  
 387 Caucasian athletes failed to reach significance.<sup>10</sup> Due to the high percentage of African American  
 388 athletes in retired football player groups provides a possible understanding for the difference in  
 389 impaired fasting glucose levels between retired athletes and controls, warranting further  
 390 investigation. All studies that reported linemen to have higher impaired fasting glucose  
 391 concentration than non-linemen.<sup>5, 34, 44, 50</sup> Although the cardiovascular risks affiliated with cardio-  
 392 metabolic syndrome and increased body size are inevitable during early retirement, engagement in  
 393 physical activity during a professional sporting career may slow the progression from impaired  
 394 fasting glucose to diabetes mellitus and decreases the risk of developing atherogenic lipoprotein  
 395 profile. Initial meta-analysis of glucose identified a high level of heterogeneity ( $I^2 = 86\%$ ). Removal of  
 396 Albuquerque et al reduced heterogeneity to a moderate level ( $I^2 = 55\%$ ), indicating a significantly  
 397 lower level of fasting glucose for retired players. However, insufficient methodological information  
 398 prevents investigation into possible causes.

399 Cardio-metabolic syndrome, an established risk factor for CVD<sup>62</sup> was shown to be highly prevalent  
 400 amongst retired athletes.<sup>38, 50, 60, 63</sup> Lineman position was associated with, and in some studies  
 401 double the prevalence compared to non-linemen and comparators.<sup>5, 34, 50</sup> Three components of  
 402 cardio-metabolic syndrome; BMI  $\geq 30$ , increased impaired fasting glucose and decreased HDL, were  
 403 significantly more prevalent in linemen.<sup>50</sup> This is further supported by an association between  
 404 cardio-metabolic syndrome and increased weight gain.<sup>10, 60</sup> Athletes playing in lineman position are  
 405 exposed to increased likelihood of the development of cardio-metabolic syndrome after retirement.  
 406 However, cardio-metabolic syndrome classification may overestimate the cardiovascular risk in  
 407 larger retired players, as previously discussed; BMI is a poor indicator of body composition in this  
 408 cohort.<sup>58</sup> The average age of retired athletes in this review was 57.1 years, therefore it is  
 409 debateable how long into retirement BMI remains an inapplicable measure due to prior elite

athleticism. Identification of the high prevalence in retired athletes is important, as many of the components of cardio-metabolic syndrome are reversible with lifestyle changes; physical activity and diet. Limited data suggests that obstructive sleep apnoea maybe more prevalent after retirement, possibly explained by the previously discussed elevated BMI and increased prevalence of obesity.<sup>1, 35, 44</sup> Retired soccer players lower mean BMI of 27kg.m<sup>2</sup> was associated with a lower incidence of obstructive sleep apnoea compared to controls and retired American football players.<sup>60</sup> However, Luyster reported that despite similar levels of obesity, high sleep apnoea risk was twice that for retired football players, compared to controls, giving plausibility to other possible causes for increased prevalence, beyond BMI. As a measure of subclinical atherosclerosis, CAP and CAC are strongly and independently associated with adverse cardiovascular events.<sup>6, 84</sup> Despite high levels of physical activity throughout athletic careers, following retirement, former athletes have a prevalence and distribution of subclinical atherosclerosis similar to the general population matched for age, sex and BMI.<sup>8, 11, 34, 44, 45</sup> The presence of CAP and CAC is a sign of advanced atherosclerosis and has significant diagnostic implications. The prevalence of CAC was consistently reported in at least one third of retired players.<sup>10, 34, 44, 63</sup> CAC < 100 was present in 76% of retired players, posing concern;<sup>10</sup> as risk of an adverse cardiovascular event increases several fold higher with CAC >100.<sup>12</sup> Retired linemen are more likely to have moderate to severe presence of CAC and less likely to have an absence of CAC compared to non-linemen.<sup>5, 34</sup> Possible explanations for higher risk of moderate to severe CAC include; increased prevalence of obesity, hypertension, cardio-metabolic syndrome and sleep-disordered breathing. However, it is difficult to rule out factors beyond those measured in these studies, including but not limited to; steroid use, race and socio-economic status. These findings suggest former athletes have not benefited from their athletic pasts, despite the well documented cardio-protective benefits associated with pro-longed engagement in exercise.<sup>54</sup>

This review is limited by several factors. First, studies included were cross sectional, observational or prospective; therefore, inferences on temporality and causality cannot be made from the observed findings. Results should be viewed as hypothesis generating only. Second, 12 of

the 13 studies included were based on retired football players; therefore, caution is needed when interpreting conclusions to all retired field-based athletes. It's worth noting that all studies included male athletes, retired from professional sports, therefore applicability to amateur and female athletes is limited. A high proportion of retired athletes, primarily football players, were African-American, limiting generalisability of results. Caution is needed when interpreting findings given the disproportionate percentage of African-American retired football players, who have a higher pre-disposition for increased BP and hypertension.<sup>46</sup> Due to the limited amount of research in this area the inclusion of a controls group was not implemented into our inclusion criteria as to widen the number of studies that could be analysed. All studies recruited participants from open health screening events allowing for self-selection bias; however, this applies to all participants; linemen, non-linemen and retired soccer players. There are possible unknown causes for findings, such as; previous use of anabolic androgenic steroids could have multiple deleterious effects on cardiovascular system; altering lipid profiles, promoting atherosclerosis, enhancing thrombogenesis and altering body composition. Other possible co-founding factors include; years in retirement, diet, alcohol use, socioeconomic status, education, genetics and medication use. Use of self-reported screening tools for obstructive sleep apnoea without objective assessment, precludes confirmation that individuals who scored in the high-risk range actually had obstructive sleep apnoea. Therefore, the proportion of high-risk participants may be overestimated or underestimated. Finally, it's difficult to acquire a similar comparator population. General population and cohorts from larger studies for the most were used; however, not all studies matched controls for ethnicity, race and body composition.

457

## 458 **Conclusion**

459 There is inconsistency in the screening and reporting of CVD risk factors in retired field-based  
460 athletes. Most studies have focused on retired football players, with only one study examining

461 retired soccer players. There is a need for research in field-based athletes from other sports,  
462 particularly in sports which emphasises greater player mass. There is also a need for research on  
463 cardiovascular risks in female athletes from similar sports. This current synthesis of studies has  
464 demonstrated that heavier retired field-based athletes are at a risk of elevated BP, hypertension,  
465 increased LDL, sleep-disordered breathing, cardio-metabolic syndrome and development of CAP and  
466 CAC. It can be inferred that this risk is comparable to obese non-athlete counterparts. BMI might not  
467 be an appropriate measurement of cardiovascular health in retired field-based athletes, and other  
468 measures of body composition may be more valuable. Further research is needed, focusing on  
469 retired athletes of other field-based sports such as rugby, hockey and soccer, and retired women  
470 athletes, to gain a clear insight into the cardiovascular health of all field-based sports.

471

472

473

474

475

476

477

478

479

480

481

482

## 483 References

- 484 1. Albuquerque FN, Kuniyoshi FH, Calvin AD, et al. Sleep-disordered breathing, hypertension,  
485 and obesity in retired National Football League players. *J Am Coll Cardiol.* 2010;56(17):1432-  
486 1433.
- 487 2. Anzell AR, Potteiger JA, Kraemer WJ, Otieno S. Changes in height, body weight, and body  
488 composition in American football players from 1942 to 2011. *J Strength Cond Res.*  
489 2013;27(2):277-284.
- 490 3. Barlow et al. Percision error in dual energy X-ray absorptiometry body composition  
491 measurements in elite male rugby players. 2015.
- 492 4. Baron SR, R. . *NIOSH mortality study of NFL football players: 1959-1988.* Cincinnati, OH:  
493 National Institute of Occupational Safety and Health; 1994.
- 494 5. Basra SS, Pokharel Y, Hira RS, et al. Relation between playing position and coronary artery  
495 calcium scores in retired National Football League players. *Am J Cardiol.* 2014;114(12):1836-  
496 1840.
- 497 6. Belcaro G, Nicolaides AN, Ramaswami G, et al. Carotid and femoral ultrasound morphology  
498 screening and cardiovascular events in low risk subjects: a 10-year follow-up study (the  
499 CAFES-CAVE study1). *Atherosclerosis.* 2001;156(2):379-387.
- 500 7. Brignole M. 'Ten Commandments' of ESC Syncope Guidelines 2018: The new European  
501 Society of Cardiology (ESC) Clinical Practice Guidelines for the diagnosis and management of  
502 syncope were launched 19 March 2018 at EHRA 2018 in Barcelona. *Eur Heart J.*  
503 2018;39(21):1870-1871.
- 504 8. Carruthers D, Joshi P, Maroules C, et al. Pooled cohort equation and coronary artery calcium  
505 in retired national football league athletes. *Journal of the American College of Cardiology.*  
506 2017;69(11):1449.
- 507 9. Caselli S, Di Paolo FM, Pisicchio C, Pandian NG, Pelliccia A. Patterns of left ventricular  
508 diastolic function in Olympic athletes. *J Am Soc Echocardiogr.* 2015;28(2):236-244.
- 509 10. Chang AY, FitzGerald SJ, Cannaday J, et al. Cardiovascular Risk Factors and Coronary  
510 Atherosclerosis in Retired National Football League Players. *American Journal of Cardiology.*  
511 2009;104(6):805-811.
- 512 12. Church TS, Levine BD, McGuire DK, et al. Coronary artery calcium score, risk factors, and  
513 incident coronary heart disease events. . *Atherosclerosis.* 2007;190(1):224-231.
- 514 13. Croft LB, Belanger A, Miller MA, Roberts A, Goldman ME. Comparison of National Football  
515 League Linemen Versus Nonlinemen of Left Ventricular Mass and Left Atrial Size. *American*  
516 *Journal of Cardiology.* 2008;102(3):343-347.
- 517 14. De Backer G, Ambrosioni E, Borch-Johnsen K, et al. European guidelines on cardiovascular  
518 disease prevention in clinical practice. Third Joint Task Force Of European and other societies  
519 on cardiovascular disease prevention in clinical practice (constituted by representatives of  
520 eight societies and by invited experts). *Arch Mal Coeur Vaiss.* 2004;97(10):1019-1030.
- 521 15. Denke MA, Sempos CT, Grundy SM. Excess body weight: an underrecognized contributor to  
522 high blood cholesterol levels in white American men. *Archives of internal medicine.*  
523 1993;153(9):1093-1103.
- 524 16. DiCesare CA, Adams JR, Claytor RP, Ward RM, Cox RH. Relationship between proxies for Type  
525 II fiber type and resting blood pressure in Division I American Football Athletes. *International*  
526 *Journal of Health Sciences-Ijhs.* 2017;11(2).
- 527 17. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical appraisal tool to  
528 assess the quality of cross-sectional studies (AXIS). *BMJ Open.* 2016;6(12):e011458.
- 529 18. European Society of Hypertension. European Society of Cardiology guidelines for the  
530 management of arterial hypertension. *Journal of hypertension.* 2003;21(6):1011-1053.
- 531 19. Fontana L, Eagon JC, Trujillo ME, Scherer PE, Klein S. Visceral fat adipokine secretion is  
532 associated with systemic inflammation in obese humans. *Diabetes.* 2007;56(4):1010-1013.



- 533 20. Ford ES. Risks for all-cause mortality, cardiovascular disease, and diabetes associated with  
534 the metabolic syndrome: a summary of the evidence. *Diabetes care*. 2005;28(7):1769-1778.
- 535 21. Fox CS, Pencina MJ, Wilson PW, Paynter NP, Vasan RS, D'agostino RB. Lifetime risk of  
536 cardiovascular disease among individuals with and without diabetes stratified by obesity  
537 status in the Framingham heart study. *Diabetes care*. 2008.
- 538 22. Fryar CD, Gu Q, Ogden CL, Flegal KM. Anthropometric Reference Data for Children and  
539 Adults: United States, 2011-2014. *Centers for Disease Control*. 2016 ed. United States:  
540 Centers for Disease and Control; 2016.
- 541 23. Garry JP, McShane JJ. Analysis of lipoproteins and body mass index in professional football  
542 players. *Prev Cardiol*. 2001;4(3):103-108.
- 543 24. Georgeson EC, Weeks BK, McLellan C, Beck BR. Seasonal change in bone, muscle and fat in  
544 professional rugby league players and its relationship to injury: a cohort study. *BMJ Open*.  
545 2012;2(6).
- 546 25. Gibbons GH, Shurin SB, Mensah GA, Lauer MS. Refocusing the agenda on cardiovascular  
547 guidelines: an announcement from the National Heart, Lung, and Blood Institute. *J Am Coll*  
548 *Cardiol*. 2013;62(15):1396-1398.
- 549 26. Greenland P, Alpert JS, Beller GA, et al. 2010 ACCF/AHA guideline for assessment of  
550 cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology  
551 Foundation/American Heart Association task force on practice guidelines developed in  
552 collaboration with the American Society of Echocardiography, American Society of Nuclear  
553 Cardiology, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular  
554 Angiography and Interventions, Society of Cardiovascular Computed Tomography, and  
555 Society for Cardiovascular Magnetic Resonance. *Journal of the American College of*  
556 *Cardiology*. 2010;56(25):e50-e103.
- 557 27. Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the metabolic  
558 syndrome: an American Heart Association/National Heart, Lung, and Blood Institute  
559 scientific statement: Executive Summary. *Crit Pathw Cardiol*. 2005;4(4):198-203.
- 560 28. Hargrove T. Supersized in the NFL: many ex-players dying young. Scripps Howard News  
561 Service Online: Scripps Howard News Service Online; 2007.
- 562 29. Haskell WL, Lee I-M, Pate RR, et al. Physical activity and public health: updated  
563 recommendation for adults from the American College of Sports Medicine and the American  
564 Heart Association. *Circulation*. 2007;116(9):1081.
- 565 30. Helzberg JH, Waeckerle JF, Camilo J, et al. Comparison of cardiovascular and metabolic risk  
566 factors in professional baseball players versus professional football players. *American*  
567 *Journal of Cardiology*. 2010;106(5):664-667.
- 568 31. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in*  
569 *medicine*. 2002;21(11):1539-1558.
- 570 32. Hu FB, Willett WC, Li T, Stampfer MJ, Colditz GA, Manson JE. Adiposity as compared with  
571 physical activity in predicting mortality among women. *N Engl J Med*. 2004;351(26):2694-  
572 2703.
- 573 33. Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for  
574 cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study.  
575 *Circulation*. 1983;67(5):968-977.
- 576 34. Hurst RT, Burke RF, Wissner E, et al. Incidence of Subclinical Atherosclerosis as a Marker of  
577 Cardiovascular Risk in Retired Professional Football Players. *American Journal of Cardiology*.  
578 2010;105(8):1107-1111.
- 579 35. Hyman MH, Dang DL, Liu Y. Differences in obesity measures and selected comorbidities in  
580 former National Football League professional athletes. *J Occup Environ Med*.  
581 2012;54(7):816-819.
- 582 36. Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management  
583 of overweight and obesity in adults: a report of the American College of

Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Journal of the American college of cardiology*. 2014;63(25 Part B):2985-3023.

37. Jones B, Till K, Barlow M, Lees M, O'Hara JP, Hind K. Anthropometric and Three-Compartment Body Composition Differences between Super League and Championship Rugby League Players: Considerations for the 2015 Season and Beyond. *PLoS One*. 2015;10(7):e0133188.

38. Kelly DF, Chaloner C, Evans D, et al. Prevalence of pituitary hormone dysfunction, metabolic syndrome, and impaired quality of life in retired professional football players: a prospective study. *J Neurotrauma*. 2014;31(13):1161-1171.

39. Kriska AMaC, C.J. Introduction to the collection of physical activity questionnaires. *Med Sci Sports Exerc*. 1997;29(6):5-9.

40. Kujala UM, Kaprio J, Taimela S, Sarna S. Prevalence of diabetes, hypertension, and ischemic heart disease in former elite athletes. *Metabolism-Clinical and Experimental*. 1994;43(10):1255-1260.

41. Lee CD, Blair SN, Jackson AS. Cardiorespiratory fitness, body composition, and all-cause and cardiovascular disease mortality in men. *Am J Clin Nutr*. 1999;69(3):373-380.

42. Lees MJ, Oldroyd B, Jones B, et al. Three-Compartment Body Composition Changes in Professional Rugby Union Players Over One Competitive Season: A Team and Individualized Approach. *J Clin Densitom*. 2017;20(1):50-57.

43. Levine DA, Lewis CE, Williams OD, et al. Geographic and demographic variability in 20-year hypertension incidence: the CARDIA study. *Hypertension*. 2011;57(1):39-47.

44. Luyster FS, Dunn RE, Lauderdale DS, et al. Sleep-apnea risk and subclinical atherosclerosis in early-middle-aged retired National Football League players. *Nature and Science of Sleep*. 2017;9:31-38.

45. Lynch NA, Ryan AS, Evans J, Katzell LI, Goldberg AP. Older elite football players have reduced cardiac and osteoporosis risk factors. *Medicine & Science in Sports & Exercise*. 2007;39(7):1124-1130.

47. Mancia G, Fagard R, Narkiewicz K, et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Journal of Hypertension*. 2013;31(7):1281-1357.

48. Mathews DK, Fox EL, Tanzi D. Physiological responses during exercise and recovery in a football uniform. *J Appl Physiol*. 1969;26(5):611-615.

50. Miller MA, Croft LB, Belanger AR, et al. Prevalence of Metabolic Syndrome in Retired National Football League Players. *American Journal of Cardiology*. 2008;101(9):1281-1284.

51. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009;6(7):p.e1000097.

52. Morehen JC, Routledge HE, Twist C, Morton JP, Close GL. Position specific differences in the anthropometric characteristics of elite European Super League rugby players. *Eur J Sport Sci*. 2015;15(6):523-529.

53. Muntner P, Lewis CE, Diaz KM, et al. Racial differences in abnormal ambulatory blood pressure monitoring measures: Results from the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Am J Hypertens*. 2015;28(5):640-648.

54. Myers J, Prakash M, Froelicher V, Do D, Partington S, Edwin Atwood J. Exercise capacity and mortality among men referred for exercise testing. *N Eng J of Med*. 2002;346(11):793-801.

55. Nevill AM, Stewart AD, Olds T, Holder R. Relationship between adiposity and body size reveals limitations of BMI. *Am J Phys Anthropol*. 2006;129(1):151-156.

56. Nevill AM, Winter EM, Ingham S, Watts A, Metsios GS, Stewart AD. Adjusting athletes' body mass index to better reflect adiposity in epidemiological research. *J Sports Sci*. 2010;28(9):1009-1016.

57. Newman AB, Foster G, Givelber R, Nieto FJ, Redline S, Young T. Progression and regression of sleep-disordered breathing with changes in weight: the Sleep Heart Health Study. *Arch Intern Med.* 2005;165(20):2408-2413.
58. Ode JJ, Pivarnik JM, Reeves MJ, Knous JL. Body mass index as a predictor of percent fat in college athletes and nonathletes. *Med Sci Sports Exerc.* 2007;39(3):403-409.
59. Paffenbarger RS, Hyde RT, Wing AL, Steinmetz CH. A natural history of athleticism and cardiovascular health. *Jama.* 1984;252(4):491-495.
60. Panayiotoglou A, Grammatikopoulou MG, Maraki MI, et al. Metabolic syndrome in retired soccer players: A pilot study. *Obesity Medicine.* 2017;8:15-22.
61. Park HS, Park JY, Yu R. Relationship of obesity and visceral adiposity with serum concentrations of CRP, TNF- $\alpha$  and IL-6. *Diabetes research and clinical practice.* 2005;69(1):29-35.
62. Perk J, De Backer G, Gohlke H, et al. European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). *Eur Heart J.* 2012;33(13):1635-1701.
63. Pokharel Y, Macedo FY, Nambi V, et al. Neck circumference is not associated with subclinical atherosclerosis in retired national football league players. *Clinical Cardiology.* 2014;37(7):402-407.
64. Punjabi NM, Caffo BS, Goodwin JL, et al. Sleep-disordered breathing and mortality: a prospective cohort study. *PLoS medicine.* 2009;6(8):e1000132.
65. Rabkin SW, Leiter L, Reeder BA, Chen Y, Liu L. Risk factor correlates of body mass index. *Canadian Medical Association. Journal.* 1997;157(1):S26.
66. Reiner Z, Catapano AL, De Backer G, et al. ESC/EAS Guidelines for the management of dyslipidaemias: the Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *European heart journal.* 2011;32(14):1769-1818.
67. Ryden L, Standl E, Bartnik M, et al. Guidelines on diabetes, pre-diabetes, and cardiovascular diseases: executive summary. The Task Force on Diabetes and Cardiovascular Diseases of the European Society of Cardiology (ESC) and of the European Association for the Study of Diabetes (EASD). *Eur Heart J.* 2007;28(1):88-136.
68. Selden MA, Helzberg JH, Waeckerle JF, et al. Cardiometabolic Abnormalities in Current National Football League Players. *American Journal of Cardiology.* 2009;103(7):969-971.
69. Shaw LJ, Raggi P, Schisterman E, Berman DS, Callister TQ. Prognostic value of cardiac risk factors and coronary artery calcium screening for all-cause mortality. *Radiology.* 2003;228(3):826-833.
70. Shea JL, Randell EW, Sun G. The prevalence of metabolically healthy obese subjects defined by BMI and dual-energy X-ray absorptiometry. *Obesity (Silver Spring).* 2011;19(3):624-630.
71. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *The Journal of clinical investigation.* 2006;116(7):1793-1801.
72. Silva GE, Vana KD, Goodwin JL, Sherrill DL, Quan SF. Identification of patients with sleep disordered breathing: comparing the four-variable screening tool, STOP, STOP-Bang, and Epworth Sleepiness Scales. *J Clin Sleep Med.* 2011;7(5):467-472.
73. Smith SC, Allen J, Blair SN, et al. AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update: endorsed by the National Heart, Lung, and Blood Institute. *Journal of the American College of Cardiology.* 2006;47(10):2130-2139.
74. Somers VK, White DP, Amin R, et al. Sleep apnea and cardiovascular disease: An american heart association/american college of cardiology foundation scientific statement from the american heart association council for high blood pressure research professional education

committee, council on clinical cardiology, stroke council, and council on cardiovascular nursing in collaboration with the national heart, lung, and blood institute national center on sleep disorders research (national institutes of health). *Journal of the American College of Cardiology*. 2008;52(8):686-717.

75. Steffes GD, Megura AE, Adams J, et al. Prevalence of metabolic syndrome risk factors in high school and NCAA division I football players. *Journal of Strength and Conditioning Research*. 2013;27(7):1749-1757.

76. Stein JH, Korcarz CE, Hurst RT, et al. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force endorsed by the Society for Vascular Medicine. *Journal of the American Society of Echocardiography*. 2008;21(2):93-111.

77. Tucker AM, Vogel RA, Lincoln AE, et al. Prevalence of cardiovascular disease risk factors among national football league players. *JAMA - Journal of the American Medical Association*. 2009;301(20):2111-2119.

78. Virani SS, Pompeii L, Lincoln AE, et al. Association between traditional cholesterol parameters, lipoprotein particle concentration, novel biomarkers and carotid plaques in retired National Football League players. *Atherosclerosis*. 2012;222(2):551-556.

79. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Int J Surg*. 2014;12(12):1495-1499.

80. Warren JM, Ekelund U, Besson H, Mezzani A, Geladas N, Vanhees L. Assessment of physical activity—a review of methodologies with reference to epidemiological research: a report of the exercise physiology section of the European Association of Cardiovascular Prevention and Rehabilitation. *European Journal of Cardiovascular Prevention & Rehabilitation*. 2010;17(2):127-139.

81. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2018;71(19):e127-e248.

82. Wolf-Maier K, Cooper RS, Banegas JR, et al. Hypertension prevalence and blood pressure levels in 6 European countries, Canada, and the United States. *Jama*. 2003;289(18):2363-2369.

83. Wright CJ, Abbey EL, Brandon BA, Reisman EJ, Kirkpatrick CM. Cardiovascular disease risk profile of NCAA Division III intercollegiate football athletes: a pilot study. *Physician and Sportsmedicine*. 2017;45(3):280-285.

84. Wyman RA, Mays ME, McBride PE, Stein JH. Ultrasound-detected carotid plaque as a predictor of cardiovascular events. *Vasc Med*. 2006;11(2):123-130.

85. Yamamoto JB, Yamamoto BE, Yamamoto PP, Yamamoto LG. Epidemiology of college athlete sizes, 1950s to current. *Research in Sports Medicine*. 2008;16(2):111-127.

86. Yudkin JS, Kumari M, Humphries SE, Mohamed-Ali V. Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? *Atherosclerosis*. 2000;148(2):209-214.

87. Zalesin KC, Franklin BA, Miller WM, Peterson ED, McCullough PA. Impact of obesity on cardiovascular disease. *Medical Clinics*. 2011;95(5):919-937.

## Appendix 1: Study Details

Lead Author (Year)	Study Design	Primary Aims	Setting	Participants	Variables	Risk Factor Prevalence
Miller <sup>50</sup> (2008)	Cross-sectional prevalence	Assess the prevalence of CMS in athletes. Assess the prevalence of CMS based on playing position.	Living heart foundation-health screening, 2004-2006	NFL, n=510 Males Mean Age: 53.8 Sex matched controls from NHANES study LM v NLM	BMI, BF%, SBP, DBP, HT, TC, TG, HDL, LDL, IFG, CMS, Smoking, DM	CMS more prevalent in LM than NLM (59.8% v 30.1%, p=0.001). Elevated BMI, IFG and reduced HDL was more prevalent in LM than NLM.
Panayiotoglou <sup>60</sup> (2017)	Cross-sectional case-control	To determine the risk and prevalence of CMS in retired professional soccer players.	Greece	Soccer, n=12 Males Mean Age: 46.7 Age-sex-BMI matched nonathletic controls	BMI, BF%, WHR, BP, snoring, smoking, TG, TC/HDL, nonHDL/HDL	Prevalence of CMS was indifferent between groups. Retired players with CMS gained significantly more weight since retirement.
Barsa <sup>5</sup> (2014)	Cross-sectional	Evaluate the presence and severity of subclinical atherosclerosis. Evaluate whether linemen position is independently associated with an increased risk of subclinical atherosclerosis.	Living heart foundation-health screening, 2007-2009	NFL, n=931 Males Mean Age: 54 No comparators LM v NLM	BMI, WC, SBP, hsCRP, TC, HDL-C, LDL-C, TG, FG, CMS, HT, DM, Smoking, CAC	LM less likely to have absence of CAC (33.8% v 41.7%, p=0.02) and greater likelihood of moderate to severe CAC (32.9% v 26.4%, p=0.04).
Chang <sup>10</sup> (2009)	Cross-sectional	Assess the prevalence of CAC in retired NFL players compared to physically active preventive medicine controls.	Living heart foundation-health screening, 2007	NFL, n=201 Males Mean Age: 51.2	BMI, WC, WHR, SBP, DBP, Fasting insulin, FG, TC, HDL,	No significant difference in the prevalence of CAC (46% v 48.3%, p=0.69) or distribution (p=0.11)

**Commented [AK24]:** Query 11: Confirm edits to this column for conciseness. Make any further edits to reduce the overall width of the table.

**Commented [U25R24]:** Edits confirmed. Amendments made to reduce width of table.

		Evaluate retired players true risk of an adverse CV event.		Age - sex- BMI- ethnicity matched participants from the DHS and ACLS	LDL, TG, HbA1C, CMS, DM, Smoking, CAC, HT	between retired players and controls.
Hurst <sup>34</sup> (2010)	Cross-sectional	Evaluate subclinical atherosclerosis in retired NFL players. Assess CV risk in professional football players.	Living heart foundation-health screening, 2006-2007	NFL, n=201 Males Mean Age: 50.8 Age-sex- BMI- smoking prevalence matched controls from Mayo clinic, 2006-2007 LM v NLM	BMI, Smoking, HT, SBP, DBP, HL, TC, HDL, LDL, TG, TC/HDL, Glucose, CAP.	Prevalence of CAP in players not significantly different to BMI matched controls (33.3% v 29.3%, p=0.45). CMS more prevalent in LM than NLM (45.8% v 22.5%, p=0.001).
Hyman <sup>35</sup> (2012)	Observational	Validate the accuracy of BMI when measuring obesity in retired NFL population. Investigate the correlation between obesity and several co-morbidities in this population.	An internal medicine practice, May 2010- June 2011	NFL, n=129 Males Mean Age: 42.2 No comparators LM v NLM	BMI, HT, OSA, LVH, DM.	BMI has poor specificity (0.36) in classifying obesity in retired football players. BMI-obesity was correlated to LM (p<0.0001) and OSA (p= 0.0005).
Albuquerque <sup>1</sup> (2010)	Cross-sectional	Assess the prevalence of SDB and HT. Compare the risk of CVD between retired NFL players and controls.	Living heart foundation-health screening, 2006	NFL, n=257 Males Mean Age: 53.9 Age-sex-BMI- matched cohort from the NHANES study	BMI, SBP, DBP, HT, Obesity, TC, TG, HDL, LDL, FG, DM, Smoking, Apnoea - hypopnoea, SDB.	SDB was present in 52.3% of retired players. The prevalence of HT and obesity (p<0.001) were higher in retired players. LM were more likely to have SDB (61.3% Vs 46.6%, p=0.02) and obesity (83.5% Vs 52.5%, p<0.001) compared with NLM. Retired players had lower TC, TG, HDL and IFG than controls.

Carruthers <sup>8</sup> (2017)	Cross-sectional	Assess the 10-y risk of atherosclerotic CVD in elite former athletes.	Not specified	NFL, n=104 Males Mean Age: 53.8 Age - sex matched participants from the DHS	BMI, SBP, Non-HDL, HDL, Median CAC, median ASCVD risk, Smoking.	No significant differences in the odds of having CAC=0 among high ASCVD risk participants (OR 1.37; 95% CI: 0.36, 5.17) nor in the odds of having high CAC (CAC>100) among low ASCVD risk participants (OR 1.28; 95% CI: 0.64, 2.54).
Kelly <sup>38</sup> (2014)	Prospective	Determine the rate of metabolic dysfunction in retired NFL players.	St. John's Health Center, Santa Monica. LA Biomedical Research Institute at Harbour-UCLA medical center.	NFL, n=74 Males Mean Age: 47.3 No comparators Non-hormone deficient v Hormone deficient	BMI, CMS, IGF-1	CMS was present in 50% of retired players. BMI increased significantly (p<0.001) for players during retirement.
Lynch <sup>45</sup> (2007)	Cross-sectional	Determine if playing professional football as a young adult is associated with a more favourable CV risk profile, and a greater bone density and lean body mass compared with their healthy peers.	University of Maryland	NFL, n=16 Males Mean Age: 66 Sex -BMI-race and current physical activity matched never-athletic comparators.	BMI, WC, WHR, BF%, TC, LDL, HDL, TG, Fasting Insulin, Fasting Glucose, BP	Retired players had more favourable body composition and CV risk profile; 37% higher HDL, fourfold higher HDL2, 25% lower TC/HDL ratio, and 31% lower TG than controls (p<0.05 to <0.001).
Pokharel <sup>63</sup> (2014)	Cross-sectional	Examine the association of NC with other markers of adiposity and components of MS. Examine whether NC was independently associated with subclinical atherosclerosis as assessed by CAC and CAP.	Player Care Foundation and the Living Heart Foundation and Boon Heart Institute	NFL, n=845 Males Mean Age: 54 No comparators	HT, DM, SBP, DBP, BMI, NC, WC, FBG, hsCRP, TC, LDL, HDL, TG, CMS, CAC/CAP	21% had CMS, 62% had CAC and 56% CAP present. NC was not associated with CAC or CAP after adjusting for age, race and cardio-metabolic risk factors.

Virani <sup>78</sup> (2012)	Cross-sectional	Assess whether LDL-P concentration and hsCRP can identify subclinical atherosclerosis better than traditional cholesterol parameters. Assess if hsCRP is associated with CAP in retired NFL players.	Living Heart Foundation and the Boone Heart Institute, September 2007-November 2009	NFL, n=948 Males Mean Age: 53.5 No comparators CMS v No CMS	HT, DM, MS, WC, TC, LDL, nonHDL, TG, LDL-P, HDL, hsCRP	CAP common in retired players (41%) and strongly associated with LDL-P (OR, 3.71, 95%CI: 1.16-11.84). 19.7% of retired players had CMS. Hs-CRP was not associated with carotid plaques (OR: 1.13, 95%CI 0.71-1.79).
Luyster <sup>44</sup> (2017)	Cross-sectional	Compare SA risk in early-middle-aged retired NFL players to a community cohort. Compare risk of SA based on playing position.	Player Care Foundation Cardiovascular Health Screening Program, 2007-2012	NFL, n=122 Males Mean age: 45.3 Age-sex-race-BMI cohort from the CARDIA sleep study	Smoking, WC, BMI, Obese, SBP, DBP, TC, HDL, LDL, TG, DM, FG, Sleep duration, SA risk, CAC	Retired players have a greater prevalence of high SA risk (27% v 11.5%, p=0.002) but similar prevalence of CAC as matched controls (30% v 30%, p=1).

**Abbreviations:** ASCVD - atherosclerotic cardiovascular disease; BF%- body fat percentage; BMI- body mass index; ; CAC- carotid artery calcium; CAP- carotid artery plaque; CMS- cardio-metabolic syndrome; DBP- diastolic blood pressure; DM - diabetes mellitus; HDL- high-density lipoprotein; hsCRP- high sensitive C reactive protein; HL- hyperlipidemia; HT- hypertension; IFG- impaired fasting glucose; LM - linemen; LDL- low- density lipoprotein; LDL-P- low-density lipoprotein particle number; NFL- National Football League; NLM - non-linemen; OSA – Obstructive Sleep Apnea; SA – sleep apnoea; SBP- systolic blood pressure; TC- total cholesterol; TG-triglycerides; WC- waist circumference; WHR- waist-hip-ratio.